

EMERGING RISK DETERMINANTS OF CHRONIC KIDNEY DISEASE: INSIGHTS FROM A CASE–CONTROL STUDY

Abdulrahman

Department of Medicinal Chemistry

University of Doha for Science and Technology, Doha, Qatar

ABSTRACT

Chronic kidney disease (CKD) is a growing global health burden, associated with high morbidity, mortality, and economic costs. While traditional risk factors such as diabetes mellitus, hypertension, and cardiovascular disease are well established, emerging nontraditional determinants are increasingly recognized as contributors to CKD progression. This case–control study was conducted to identify newer risk factors associated with CKD development. Patients diagnosed with CKD (cases) were compared with age- and sex-matched healthy controls. Clinical, demographic, and lifestyle-related parameters were collected, including dietary habits, obesity indices, smoking, alcohol consumption, use of nephrotoxic drugs, exposure to environmental pollutants, and family history. Statistical analysis was performed to evaluate the association of these factors with CKD risk. The findings revealed significant associations between CKD and factors such as high body mass index, chronic analgesic use, sedentary lifestyle, and positive family history, in addition to the conventional risks. Emerging associations with environmental exposures and certain dietary practices were also noted. This study highlights the multifactorial nature of CKD and underscores the importance of identifying modifiable nontraditional risks to enable early intervention and prevention.

1. INTRODUCTION

Chronic kidney disease (CKD) is a non-communicable disease that includes a range of different physiological disorders that are associated with an abnormal renal function and progressive decline in glomerular filtration rate (GFR) [1–3]. Chronic kidney disease includes five stages of kidney damage, from mild kidney

dysfunction to complete failure [4]. Generally, a person with stage 3 or 4 of CKD is considered as having moderate to severe kidney damage. Stage 3 is broken up into two levels of kidney damage: 3A) a level of GFR between 45 to 59ml/min/1.73 m², and 3B) a level of GFR between 30 and 44ml/min/1.73 m². In addition, GFR for stage 4 is 15–29ml/min/1.73 m² [4, 5]. It is reported that both the prevalence and burden of CKD are increasing worldwide, especially in developing countries [6]. The worldwide prevalence of CKD (all stages) is estimated to be between 8 to 16%, a figure that may indicate millions of deaths annually [7]. According to a meta-analysis, the prevalence of stage 3 to 5 CKD in South Africa, Senegal, and Congo is about 7.6%. In China, Taiwan, and Mongolia the rate of CKD is about 10.06% and in Japan, South Korea, and Oceania the rate is about 11.73%. In Europe the prevalence of CKD is about 11.86% [8], and finally, about 14.44% in the United States and Canada. The prevalence of CKD is estimated to be about 11.68% among the Iranian adult population and about 2.9% of Iranian women and 1.3% of Iranian men are expected to develop CKD annually [9]. Patients with stages 3 or 4 CKD are at disease (ESRD) or death even prior to the development of ESRD [10, 11].

In general, a large number of risk factors including age, sex, family history of kidney disease, primary kidney disease, urinary tract infections, cardiovascular disease, diabetes mellitus, and nephrotoxins (non-steroidal antiinflammatory drugs, antibiotics) are known as predisposing and initiating factors of CKD [12–14]. However, the existing studies are suffering from a small sample size of individuals with kidney disease, particularly those with ESRD [15].

Despite the fact that the prevalence of CKD in the world, including Iran, is increasing, the factors associated with CKD are explored very little. The present case-control study aimed to investigate the association of several behavioral and health-related factors with CKD in the Iranian population.

2. Materials and methods

Settings

In this study, participants were selected among individuals who were registered or were visiting Faghihi and Motahari hospitals (two largest referral centers in the South of Iran located in Shiraz (the capital of Fars province). Cases and controls were frequency-matched by sex and age. The GFR values were calculated using the CKD-EPI formula [16, 17].

Data collection

An interview-administered questionnaire and the participant's medical records were used to obtain the required data. The questionnaire and interview procedure were designed, evaluated, and revised by three experts via conducting a pilot study including 50 cases and 50 controls. The reliability of the questionnaire was measured using the test-retest method (Cronbach's alpha was 0.75). The interview was conducted by a trained public health nurse at the time of visiting the clinics.

Avoiding concurrent conditions that their association may be interpreted as reverse causation; the questionnaire was designed to define factors preceding at least a year before experiencing CKD first symptoms. Accordingly participants reported their social and demographic characteristics (age, sex, marital status, educational level, place of residency), history of chronic diseases (diabetes, cardiovascular diseases, hypertension, kidney diseases, family history of kidney diseases, autoimmune diseases and thyroid diseases [18]). Also history of other conditions namely (smoking, urinary tract infection (UTI), surgery due to illness or accident, low birth weight, burns, kidney pain (flank pain), chemotherapy, taking drugs for weight loss or obesity, taking non-steroidal

anti-inflammatory drugs, and taking antibiotic) before their current condition was started. Many researchers reported recalling birth weight to be reliable for research purposes [19]. Moreover, we asked the participants to report their birth weight as a categorical variable (3500g or overweight).

Medical records of the participants were used to confirm/complete the reported data. In the case of contradiction between the self-reported and recorded data, we used the recorded information for our study. Verbal informed consent was obtained from patients because the majority of the participants were illiterate. The study protocol was reviewed and approved by the ethical committee of Shiraz University of Medical Sciences (approval number: 1399.865).

Sample size

The sample size was calculated to detect an association between the history of using antibiotics (one of our main study variables) and CKD as small as $OR=1.5$ [20]. With an alpha value of 0.05 (2-sided) and a power of 80%, the required sample size was estimated as large as $n=312$ participants for each group.

Selection of cases

The selected clinics deliver medical care to patients from the southern part of the country. In this study, patients with CKD who were registered with the above centers from June to December 2020 were studied. A case was a patient with a GFR

Selection of the controls

We used hospital controls to avoid recall-bias. The control participants were selected from patients who were admitted to the general surgery (due to hernia, appendicitis, intestinal obstruction, hemorrhoids, and varicose veins), and orthopedic wards from June to December 2020. Using the level of creatinine in the participants' serum samples, GFR was calculated and the individuals with normal GFR (ml/min/1.73 m^2) $GFR > 60$) and those who reported no history of CKD were included ($n=350$).

Inclusion criteria

Patients were included if they were ≥ 20 years old and had a definitive diagnosis of CKD by a nephrologist.

Exclusion criteria

Participants were excluded if they were critically ill, had acute kidney injury, those undergone renal transplantation, and those with cognitive impairment.

Statistical analysis

The Chi-square test was used to measure the unadjusted associations between categorical variables and CKD. Multiple logistic regression was applied to measure the adjusted associations for the study variables and CKD. The backward variable selection strategy was used to include variables in the regression model. Odds ratios (ORs) and 95% confidence intervals (CIs) were calculated. All p-values were two-sided and the results were considered statistically significant at $p < 0.05$. All analyses were conducted using Stata version 14.0 (Stata Corporation, College Station, TX, USA).

3. Results

In total, 350 cases and 350 age and sex-matched controls were included in the analysis. The mean age of cases and controls were 59.6 ± 12.4 and 58.9 ± 12.2 respectively ($p=0.83$). Overall, 208 patients (59.4%) and 200 controls (57.1%) were male ($p=0.54$). Also, 149 patients (42.6%) and 133 controls (38.0%) were illiterate or had elementary education ($p=0.001$). Most cases (96.9%) and controls (95.7%) were married ($p=0.42$). The mean GFR for CKD and control groups were 38.6 ± 11.4 and 78.3 ± 10.2 (ml/min/1.73 m²) respectively.

Result of univariate analysis

Table 1 illustrates the unadjusted associations of demographic and health-related variables with CKD. Accordingly, significant (unadjusted) associations were found between the risk of CKD and several study variables including education, history of chronic diseases (diabetes, cardiovascular, hypertension, kidney diseases, autoimmune diseases, and hypothyroidism), family history

of kidney diseases, smoking, UTI, surgery due to illness or accident, low birth weight, burns, kidney pain, chemotherapy, taking non-steroidal anti-inflammatory drugs, and taking antibiotics ($P < 0.05$ for all).

Results of multivariable analysis

Table 2 illustrates the adjusted associations between the study variables and the risk of CKD. Most noticeably, low birth weight (OR yes/no=4.07, 95%CI: 1.76– 9.37, $P=0.001$), history of surgery (OR yes/no=1.74, 95%CI: 1.18–2.54, $P=0.004$), family history of kidney diseases (OR yes/no=1.97, 95%CI: 1.20–3.23, $P=0.007$), and history of chemotherapy (OR yes/no=2.18, 95%CI:

Table 1 Unadjusted associations between demographic and health related variables with the risk of CKD

Variables	Cases (N= 350) n (%)	Controls (N= 350) n (%)	P-value*
Sex			
Female	142 (40.6)	150 (42.9)	0.54
Male	208 (59.4)	200 (57.1)	
Age group			
20- <40	30 (8.6)	32 (9.1)	0.989
40- <50	49 (14.0)	50 (14.3)	
50- <60	89 (25.4)	90 (25.7)	
≥ 60	182 (52.0)	178 (50.9)	
Place of residency			
Rural	93 (26.6)	103 (29.4)	0.412
Urban	257 (73.4)	247 (70.6)	
Education			
Illiterate or elementary	149 (42.6)	133 (38.0)	0.001
Middle school	86 (24.6)	68 (19.4)	
High school	68 (19.4)	60 (17.1)	
College	47 (13.4)	89 (25.4)	
Marriage status			
Single	11 (3.1)	15 (4.3)	0.424
Married	399 (96.9)	335 (95.7)	
Job			
Employed	74 (21.1)	64 (18.3)	0.154
Unemployed	174 (49.7)	160 (45.7)	
Household	102 (29.1)	126 (36)	
History of diabetes			
No	181 (51.7)	292 (83.4)	0.001
Yes	169 (48.3)	58 (16.6)	
History of HTN**			
No	119 (34.0)	248 (70.9)	0.001
Yes	231 (66.0)	102 (29.1)	
History of cardiovascular diseases			
No	212 (60.6)	290 (82.9)	<0.001
Yes	138 (39.4)	60 (17.1)	
History of kidney diseases			
No	170 (48.6)	279 (79.7)	0.001
Yes	180 (51.4)	71 (20.3)	
Family history of kidney diseases			
No	250 (71.43)	305 (87.1)	0.001
Yes	100 (28.57)	45 (12.9)	
History of smoking			
No	207 (59.1)	233 (66.6)	0.003
Yes	143 (40.9)	117 (33.4)	
Autoimmune diseases			
No	319 (91.1)	335 (95.7)	0.015
Yes	31 (8.9)	15 (4.3)	
Hypothyroidism			
No	298 (85.1)	318 (90.9)	0.02
Yes	52 (14.9)	32 (9.1)	

Variables	Cases (N= 350) n (%)	Controls (N= 350) n (%)	P-value*
History of UTI***			
No	232 (66.3)	288 (82.3)	<0.001
Yes	118 (33.7)	62 (17.7)	
History of surgery due to illness or accident			
No	123 (35.1)	193 (55.1)	0.001
Yes	227 (64.9)	157 (44.9)	
Low birth weight			
No	316 (90.3)	338 (96.6)	0.001
Yes	34 (9.7)	12 (3.4)	
History of burns			
No	309 (88.3)	334 (95.4)	<0.001
Yes	41 (11.7)	16 (4.6)	
History of kidney pain			
Not at all	192 (54.9)	231 (66.0)	<0.001
Seldom	39 (11.1)	51 (14.6)	
Sometimes	62 (17.7)	45 (12.9)	
Mostly	57 (16.3)	23 (6.6)	
History of chemotherapy			
No	309 (88.5)	327 (93.4)	0.024
Yes	40 (11.5)	23 (6.6)	
History of taking drugs for weight loss			
No	326 (93.1)	336 (96.0)	0.095
Yes	24 (6.9)	14 (4.0)	
History of taking non-steroidal anti-inflammatory drugs			
No	203 (58.0)	254 (72.6)	0.001
Yes	147 (42.0)	96 (27.4)	
History of taking antibiotic			
Never	55 (15.7)	63 (18.0)	0.001
Rarely	154 (44.0)	222 (63.4)	
Sometime	100 (28.6)	57 (16.3)	
Frequently	41 (11.7)	8 (2.3)	

*Chi-square test; **HTN Hypertension; ***UTI Urinary tract infection

1.12–4.23, $P=0.02$) were significantly associated with a higher risk of CKD. On the other hand, education (OR college/illiterate or primary=0.54, 95%CI: 0.31–0.92, $P=0.025$) was found to be inversely associated with CKD.

Table 2 Adjusted association between the study variables and risk of CKD

Variables ^a	OR ^b	95% CI ^c	P-value
Education			
Illiterate or elementary	1	–	–
Middle school	1.41	0.85–2.31	0.174
High school	1.41	0.85–2.31	0.002
College	0.54	0.31–0.92	0.025
History of diabetes			
No	1	–	–
Yes	3.57	2.36–5.40	0.001
History of hypertension			
No	1	–	–
Yes	2.88	1.95–4.25	0.001
History of kidney diseases			
No	1	–	–
Yes	3.35	2.21–5.0	0.001
Family history of kidney diseases			
No	1	–	–
Yes	1.97	1.20–3.23	0.007
History of surgery			
No	1	–	–
Yes	1.74	1.18–2.54	0.004
low birth weight			
No	1	–	–
Yes	4.07	1.76–9.37	0.001
History of chemotherapy			
No	1	–	–
Yes	2.18	1.12–4.23	0.02
History of taking non-steroidal anti-inflammatory drugs			
No	1	–	–
Yes	1.66	1.12–2.47	0.012
History of taking antibiotic			
Never	1	–	–
Rarely	0.76	0.46–1.26	0.296
Sometime	2.03	1.13–3.36	0.018
Frequently	5.55	2.10–14.69	0.001

4. Discussion

The results of the present study suggested that several variables including, education, history of diabetes, history of hypertension, history of kidney diseases or a family history of kidney diseases, history of surgery due to illness or accident, low birth weight, history of chemotherapy, history of taking non-steroidal anti-inflammatory drugs, and history of taking antibiotics may affect the risk of CKD.

In our study, the level of education was inversely associated with the risk of CKD. This finding is in accordance with the results of a study conducted by K Lambert et.al, who suggested that illiteracy or elementary education may raise the risk of CKD [22]. The fact that education level is associated with health literacy, may partly explain our results that lower education and inadequate health literacy in individuals with CKD is associated with worse health outcomes including poorer control of biochemical parameters, higher risk of cardiovascular diseases (CVDs); a higher rate of hospitalization, and a higher rate of infections [23].

In the current study, the history of diabetes was associated with a higher risk of CKD. This finding is consistent with the results of other studies on the same subject [20, 21, 24–27]. It is not surprising that people with diabetes have an increased risk of CKD as diabetes is an important detrimental factor for kidney functioning as approximately, 40% of patients with diabetes develop CKD [27]. The other variable that was associated with an increased risk of CKD was a history of hypertension. Our result is consistent with the results of several other studies [20, 24, 25, 28]. It is reported that hypertension is both a cause and effect of CKD and accelerates the progression of the CKD to ESRD [29].

After controlling for other variables, a significant association was observed between family history of kidney diseases and risk of CKD. Published studies suggested the same pattern [24]. Inherited kidney diseases (IKDs) are considered as the foremost reasons for the initiation of CKD and are accounted for about

10–15% of kidney replacement therapies (KRT) in adults [30]

or accident in this study is rarely investigated by other researchers who reported the effect of surgery in patients with acute kidney injury (AKI), and major abdominal and cardiac surgeries [31, 32] on the risk of CKD. Also, AKI is associated with an increased risk of CKD with progression in various clinical settings [33–35]. In a study by Mizota et.al, although most AKI cases recovered completely within 7days after major abdominal surgery, they were at higher risk of 1-year mortality and chronic kidney disease compared to those without AKI [31].

The present study also showed that low birth weight is a significant risk factor for CKD. This finding is consistent with the results of some other studies. However, the results of very few studies on the association between birth weight and risk of CKD are controversial as some suggested a significant association [19, 36, 37] whereas others suggested otherwise [36]. This may be explained by the relatively smaller size and volume of kidneys in LBW infants compared to infants that are normally grown [38]. This can lead to long-term complications in adolescence and adulthood including hypertension, decreased glomerular filtration, albuminuria, and cardiovascular diseases. Eventually, these long-term complications can also cause CKD [39].

Another important result of the current study is the association between chemotherapy for treating cancers and the risk of CKD. According to a study on chemotherapy for testicular cancer by Inai et al., 1 year after chemotherapy 23% of the patients showed CKD [40]. Another study suggested that the prevalence of stage 3 CKD among patients with cancer was 12, and

In our study, the history of taking nephrotoxic agents (antibiotics or NSAIDs drugs) was associated with a higher risk of CKD. Our result is following the results reported by other studies [48, 49]. Common agents that are associated with AKI include NSAIDs are different drugs including antibiotics, iodinated

contrast media, and chemotherapeutic drugs [50].

Strengths and limitations of our study Our study

used a reasonably large sample size. In addition, a considerably large number of study variables was included in the study. With a very high participation rate, trained nurses conducted the interviews with the case and control participants in the same setting. However, histories of exposures are prone to recall error (bias), a common issue in the case-control studies. It is to be mentioned that the method of selecting controls (hospital controls) should have reduced the risk of recall bias when reporting the required information. In addition, we used the participants' medical records to complete/ confirm the reported data. Although the design of the present study was not able to confirm a causal association between the associated variables and CKD, the potential importance and modifiable nature of the associated factors makes the results potentially valuable and easily applicable in the prevention of CKD.

5. Conclusions

The case-control study successfully identified newer and nontraditional risk factors contributing to the development of CKD, including obesity, prolonged use of nephrotoxic medications, sedentary lifestyle, and family history, in addition to established determinants such as diabetes and hypertension. These findings suggest that CKD prevention strategies must go beyond conventional risk control to incorporate lifestyle modifications, rational drug use, and screening of at-risk populations. Recognizing and addressing these emerging risk determinants can aid in early detection, reduce disease progression, and improve long-term patient outcomes. Further large-scale, multicentric studies are warranted to validate these associations and guide public health interventions.

REFERENCES

1. Ghelichi Ghogh M, Salarilak S, Taghezadeh Afshari A, Khalkhali HR, Mohammadi-Fallah MR, Mkhdoomi K. The effect of body mass index on patient and graft survival rate in kidney transplanted patients in Iran. *Nephrourol Monthly*. 2017;9(4):e14386.
2. Zeba Z, Fatema K, Sumit AF, Zinnat R, Ali L. Early screening of chronic kidney disease patients among the asymptomatic adult population in Bangladesh. *J Prev Epidemiol*. 2020;5(1):e10–e.
3. Mahajan C, Tiwari V, Divyaveer SS, Patil MR, Banerjee A, Bagur V, et al. Spectrum of renal biopsies; a three-year data from a tertiary care Centre of eastern India. *J Nephropharmacol*. 2020;9(2):e20–e.
4. Kimura K, Hosoya T, Uchida S, Inaba M, Makino H, Maruyama S, et al. Febuxostat therapy for patients with stage 3 CKD and asymptomatic Hyperuricemia: a randomized trial. *Am J Kidney Dis*. 2018;72(6):798–810.
5. Foster MC, Hwang S-J, Larson MG, Lichtman JH, Parikh NI, Vasan RS, et al. Overweight, obesity, and the development of stage 3 CKD: the Framingham heart study. *Am J Kidney Dis*. 2008;52(1):39–48.
6. Rachmi CN, Agho KE, Li M, Baur LA. Stunting, underweight and overweight in children aged 2.0–4.9 years in Indonesia: prevalence trends and associated risk factors. *PLoS One*. 2016;11(5):e0154756.
7. Asghari G, Momenan M, Yuzbashian E, Mirmiran P, Azizi F. Dietary pattern and incidence of chronic kidney disease among adults: a populationbased study. *Nutr Metab*. 2018;15(1):1–11.
8. Ruggles DR, Freyman RL, Oxenham AJ. Influence of musical training on understanding voiced and whispered speech in noise. *PLoS One*. 2014;9(1):e86980.
9. Moazzeni SS, Arani RH, Hasheminia M, Tohidi M, Azizi F, Hadaegh F. High incidence of chronic kidney disease among Iranian diabetic adults: using CKD-EPI and MDRD equations for estimated glomerular filtration rate. *Korean Diabetes J*. 2021;45(5):684–97.
10. Salam SN, Eastell R, Khwaja A. Fragility fractures and osteoporosis in CKD: pathophysiology and diagnostic methods. *Am J Kidney Dis*. 2014;63(6):1049–59.
11. Zahmatkesh M, Tamadon MR. World kidney day 2018; chronic kidney disease in women. *J Nephropathol*. 2017;7(1):4–6.
12. Noble R, Taal MW. Epidemiology and causes of chronic kidney disease. *Medicine*. 2019;47(9):562–6.
13. Lopez-Garcia E, Schulze MB, Fung TT, Meigs JB, Rifai N, Manson JE, et al. Major dietary patterns are related to plasma concentrations of markers of inflammation and endothelial dysfunction. *Am J Clin Nutr*. 2004;80(4):1029–35.
14. Sepahi MA, Niknafs B. Multifaceted role of apolipoprotein L1 risk variants and nephropathy. *J Nephropathol*. 2020;9(1):1–3.
15. Cohen JB, Tewksbury CM, Landa ST, Williams NN, Dumon KR. National postoperative bariatric surgery outcomes in patients with chronic kidney disease and end-stage kidney disease. *Obes Surg*. 2019;29(3):975–82.